arXiv:1610.08345v2 [math.CA] 6 Nov 2016

# APPROXIMATING $n$-TH DIFFERENTIABLE FUNCTIONS OF TWO VARIABLES AND MID-POINT FORMULA

MOHAMMAD W. ALOMARI

Abstract. In this work, approximations for real two variables function $f$ which has continuous partial $(n-1)$-derivatives $(n \geq 1)$ and has the $n$–th partial derivative of bounded bivariation or absolutely continuous are established. Explicit bounds for this representation are given. An approximation of a function $f$ by its mid-point formula with its error is established.

## 1. Introduction

The Trapezoid formula

$$\int_a^b f(x)dx \approx (b-a)\frac{f(a)+f(b)}{2} \tag{1.1}$$

is one of the most popular rule that approximate the definite integral $\int_a^b f(x)dx$ of a real function $f$ over an interval $[a,b]$.

In the last forty years, a great attention and valuable efforts were devoted to investigate this formula in several ways and for various type of functions. Among others, the generalized trapezoid formula (GFT)

$$\frac{1}{b-a}\left[(x-a)\,f\,(a)+(b-x)\,f\,(b)\right] \tag{1.2}$$

was expanded by Cerone et al. in [7], and eligible to be considered as one of the important works in the latest presented literature. So that instead of (1.1) one may use (1.2), i.e.,

$$\int_a^b f(t)dt \approx \frac{1}{b-a}\left[(x-a)\,f\,(a)+(b-x)\,f\,(b)\right], \qquad \forall x \in [a,b]. \tag{1.3}$$

In another approach, if $f:[a,b]\to\mathbb{R}$ is assumed to be bounded on $[a,b]$. The chord that connects its end points $A=(a,f(a))$ and $B=(b,f(b))$ has the equation

$$d_f:[a,b]\to\mathbb{R}, \quad d_f\,(x)=\frac{1}{b-a}\left[f\,(a)\,(b-x)+f\,(b)\,(x-a)\right].$$

Some error approximations for the value of the function $f(\mathbf{x})$ by the functional

$$\Phi_f\,(x):=\frac{b-x}{b-a}\cdot f\,(a)+\frac{x-a}{b-a}\cdot f\,(b)-f\,(x).$$

were introduced by Dragomir in [8], e.g., the following bounds for $\Phi_f\,(x)$ are hold:

*Date*: November 8, 2016.
2000 *Mathematics Subject Classification.* 26B30, 26B40, 26D10, 26D15.
*Key words and phrases.* Approximation, bounded bivariation, Function of two variables.

**Theorem 1.** *If $f:[a,b]\to\mathbb{R}$ is of bounded variation, then*

$$|\Phi_f(x)| \le \left(\frac{b-x}{b-a}\right)\cdot\bigvee_a^x(f)+\left(\frac{x-a}{b-a}\right)\cdot\bigvee_x^b(f)$$

$$\le \begin{cases} \left[\frac{1}{2}+\left|\frac{x-\frac{a+b}{2}}{b-a}\right|\right]\cdot\bigvee_a^b(f) \\ \left[\left(\frac{b-x}{b-a}\right)^p+\left(\frac{x-a}{b-a}\right)^p\right]^{\frac{1}{p}}\cdot\left[\left(\bigvee_a^x(f)\right)^q+\left(\bigvee_x^b(f)\right)^q\right]^{\frac{1}{q}}; \\ \qquad\qquad p>1,\ \frac{1}{p}+\frac{1}{q}=1 \\ \frac{1}{2}\bigvee_a^b(f)+\frac{1}{2}\left|\bigvee_a^x(f)-\bigvee_x^b(f)\right| \end{cases} \tag{1.4}$$

*The first inequality in (1.4) is sharp. The constant $\frac{1}{2}$ is best possible in the first and third branches.*

In 2008, Dragomir [9] provided an approximation for the function $f$ which possesses continuous derivatives up to the order $n-1$ $(n\ge 1)$ and has the $n$-th derivative of bounded variation, in terms of the chord that connects its end points $A=(a,f(a))$ and $B=(b,f(b))$ and some more terms which depend on the values of the $k$ derivatives of the function taken at the end points $a$ and $b$, where $k$ is between 1 and $n$.

The following representation of $f$ was presented by Dragomir in [9], as follows:

**Theorem 2.** *Let $I$ be a closed subinterval on $\mathbb{R}$, let $a,b\in I$ with $a<b$ and let $n$ be a nonnegative integer. If $f:I\to\mathbb{R}$ is such that the $n$-th derivative $f^{(n)}$ is of bounded variation on the interval $[a,b]$, then for any $x\in[a,b]$ we have the representation*

$$f(x)=\frac{1}{b-a}\left[(b-x)f(a)+(x-a)f(b)\right] \tag{1.5}$$
$$+\frac{(b-x)(x-a)}{b-a}\cdot\sum_{k=1}^{n}\frac{1}{k!}\left[(x-a)^{k-1}f^{(k)}(a)+(-1)^k(b-x)^{k-1}f^{(k)}(b)\right]$$
$$+\frac{1}{b-a}\int_a^b S_n(x,t)\,d\left(f^{(n)}(t)\right),$$

*where the kernel $S_n:[a,b]^2\to\mathbb{R}$ is given by*

$$S_n(x,t)=\begin{cases}(x-t)^n(b-x), & a\le t\le x \\ (-1)^{n+1}(t-x)^n(x-a), & a\le t\le x\end{cases}$$

*and the integral in the remainder is taken in the Riemann–Stieltjes sense.*

On utilizing the notations

$$D_n(f;x,a,b):=\frac{1}{b-a}\left[(b-x)f(a)+(x-a)f(b)\right]$$
$$+\frac{(b-x)(x-a)}{b-a}\cdot\sum_{k=1}^{n}\frac{1}{k!}\left[(x-a)^{k-1}f^{(k)}(a)+(-1)^k(b-x)^{k-1}f^{(k)}(b)\right] \tag{1.6}$$

with

$$E_n(f;x,a,b):=\frac{1}{b-a}\int_a^b S_n(x,t)\,d\left(f^{(n)}(t)\right), \tag{1.7}$$

and under the assumptions of Theorem 2, Dragomir provided an error approximation of the function $f$ using the formula $f(x) - D_n(f;x,a,b) = E_n(f;x,a,b)$ for any $x \in [a,b]$.

The concept of Riemann–Stieltjes ($\mathcal{RS}$) double integral $\int_a^b \int_c^d f(x,y)\, d_x d_y \alpha(x,y)$ was characterized by Fréchet in [13], and investigated later on by Clarkson in [5]. Such $\mathcal{RS}$-integral plays an important role in Mathematics with multiple applications in several subfields including Probability Theory & Statistics, Complex Analysis, Functional Analysis, Operator Theory and others.

For $a,b,c,d \in \mathbb{R}$, we consider the subset $Q := \{(x,y) : a \le x \le b, c \le y \le d\}$ of $\mathbb{R}^2$. If $P := \{(x_i,y_j) : x_{i-1} \le x \le x_i\,;\, y_{j-1} \le y \le y_j\,;\, i = 1,\dots,n\,;\, j = 1,\dots,m\}$ is a partition of $Q$, write

$$\Delta_{11} f(x_i,y_j) = f(x_{i-1},y_{j-1}) - f(x_{i-1},y_j) - f(x_i,y_{j-1}) + f(x_i,y_j)$$

for $i = 1,2,\cdots,n$ and $j = 1,2,\cdots,m$. The function $f(x,y)$ is said to be of bounded variation in the Vitali sense (or simply bounded bivariation [14]) if there exists a positive quantity $M$ such that for every partition on $Q$ we have $\sum_{i=1}^{n}\sum_{j=1}^{m} |\Delta_{11} f(x_i,y_j)| \le M$ (see [6]). Consequently, the total bivariation of a function of two variables, over bidimensional interval $Q$, is defined to be the number

$$\bigvee_Q (f) := \bigvee_c^d \bigvee_a^b (f) := \sup\left\{\sum P : P \in \mathcal{P}(Q)\right\},$$

where $\sum(P)$ denote the sum $\sum_{i=1}^{n}\sum_{j=1}^{m} |\Delta_{11} f(x_i,y_j)|$ corresponding to the partition $P$ of $Q$. For further properties of mappings of bounded bivariation we refer the interested reader to the book recent [14] and the monograph [1].

In 2009, Alomari [1] has studied the approximation problem of the Riemann–Stieltjes double integrals $\int_c^d \int_a^b f(t,s)\, d_t d_s \alpha(t,s)$ for various kind of the integrand $f$ and the integrator $\alpha$ via inequalities approach. Among others, we need the following two basic results from [1]:

**Lemma 1.** (*Integration by parts* [1]) *If $f \in \mathcal{RS}(\alpha)$ on $Q$, then $\alpha \in \mathcal{RS}(f)$ on $Q$, and we have*

$$\begin{aligned}(1.8)\quad &\int_c^d \int_a^b f(t,s)\, d_t d_s \alpha(t,s) + \int_c^d \int_a^b \alpha(t,s)\, d_t d_s f(t,s) \\ &\qquad = f(b,d)\,\alpha(b,d) - f(b,c)\,\alpha(b,c) - f(a,d)\,\alpha(a,d) + f(a,c)\,\alpha(a,c).\end{aligned}$$

**Lemma 2.** ([1]) *Assume that $g \in \mathcal{RS}(\alpha)$ on $Q$ and $\alpha$ is of bounded bivariation on $Q$, then the inequality*

$$(1.9)\qquad \left|\int_c^d \int_a^b g(x,y)\, d_x d_y \alpha(x,y)\right| \le \sup_{(x,y)\in Q} |g(x,y)| \cdot \bigvee_Q (\alpha),$$

*holds.*

A small attention and a few works have been considered for mappings of two variables; particularly, the approximation problem of $\mathcal{RS}$-double integral $\int_a^b \int_c^d f(t,s)\, d_s d_t \alpha(t,s)$

in terms of $\mathcal{RS}$-double sums. Although the important applications in approximations, numerical integration in two or more variables is still much less developed area than its one-dimensional counterpart, which has been worked on intensively. For recent inequalities involving functions of two variables the reader may refer to [1]–[4], [10]–[12] and [15]–[18].

In this paper, we extend Cerone et al [7] and Dragomir [8], [9] results (mainly Theorem 2) by giving bidimensional representation for functions whose $n$-th partial derivatives are of bounded bivariation or absolutely continuous. Explicit bounds for this representation are given. An approximation of a function $f$ by its mid-point formula with its error is established.

## 2. The Result

For a bounded function $f : Q \to \mathbb{R}$ the chord plane that connects its end points may be represented as:

$$C_f(x,y) := \frac{1}{(b-a)(d-c)}\left[(b-x)(d-y)f(a,c)+(b-x)(y-c)f(a,d)\right.$$
$$\left.+(x-a)(d-y)f(b,c)+(x-a)(y-c)f(b,d)\right],$$

which is the generalized Trapezoid formula for functions of two variables. Consequently, the difference between the function $f$ and its chord plane may considered by the functional:

$$\psi(f,C) = C_f(x,y) - f(x,y).$$

One may study this difference for various type of functions. On the other hand, this representation can be extended to $n$-th differentiable functions, as follows:

**Theorem 3.** *Let $f : Q \to \mathbb{R}$ be a real valued function which has continuous $(n-1)$ partial derivatives ($n \geq 1$). If the $n$-th partial derivatives $D^n f$ ($n \geq 1$) are of bounded bivariation on $Q$, then for any $(x,y) \in Q$ we have the representation*

$$f(x,y) = \frac{1}{(b-a)(d-c)}\left[(b-x)(d-y)f(a,c)+(b-x)(y-c)f(a,d)\right.$$
$$\left.+(x-a)(d-y)f(b,c)+(x-a)(y-c)f(b,d)\right]+\frac{(y-c)(d-y)}{(b-a)(d-c)}$$
$$\times\sum_{j=1}^{n}\frac{1}{j!}\binom{n}{j}\left\{(b-x)(x-a)^{n-j}\left[(y-c)^{j-1}D^n f(a,c)+(-1)^j(d-y)^{j-1}D^n f(a,d)\right]\right.$$
$$\left.+(x-a)(b-x)^{n-j}\left[(-1)^j(y-c)^{j-1}D^n f(b,c)+(d-y)^{j-1}D^n f(b,d)\right]\right\}$$
$$+\frac{1}{(b-a)(d-c)}\int_c^d\int_a^b S_n(x,t;y,s)\,d_t d_s\left(D^n f(t,s)\right) \tag{2.1}$$

*where, $D^n f(t,s) = \frac{\partial^n f}{\partial t^{n-j}\partial s^j}(t,s)$ $(j = 0,1,\cdots,n)$ and*

$$S_n(x,t;y,s) = \frac{1}{n!}\begin{cases}(x-t)^n(b-x)(y-s)^n(d-y), & a\leq t\leq x,\quad c\leq s\leq y\\ (-1)^n(t-x)^n(x-a)(y-s)^n(d-y), & x<t\leq b,\quad c\leq s\leq y\\ (-1)^n(x-t)^n(b-x)(s-y)^n(y-c), & a\leq t\leq x,\quad y<s\leq d\\ (t-x)^n(x-a)(s-y)^n(y-c), & x<t\leq b,\quad y<s\leq d\end{cases}$$

*Proof.* We utilize the following Taylor's representation formula for functions $f : Q \subset \mathbb{R}^2 \to \mathbb{R}$ such that the $n$-th partial derivatives $D^n f$ are of locally bounded bivariation on $Q$,

$$f(x,y) = P_n(x,y) + R_n(x,y) \tag{2.2}$$

such that,

$$P_n(x,y) = \sum_{j=0}^{n} \frac{1}{j!} \binom{n}{j} (x-x_0)^{n-j} (y-y_0)^j D^n f(x_0,y_0) \tag{2.3}$$

and

$$R_n(x,y) = \frac{1}{n!} \int_{y_0}^{y} \int_{x_0}^{x} (x-t)^n (y-s)^n \, d_t d_s \left(D^n f(t,s)\right) \tag{2.4}$$

where, $D^n f(t,s) = \frac{\partial^n f}{\partial t^{n-j} \partial s^j}(t,s)$, and $(x,y)$, $(x_0,y_0)$ are in $Q$ and the double integral in the remainder is taken in the Riemann-Stieltjes sense.

Choosing $x_0 = a$, $y_0 = c$ and then $x_0 = b$, $y_0 = d$ in (2.1) we can write that

$$\begin{aligned} f(x,y) = \sum_{j=0}^{n} \frac{1}{j!} \binom{n}{j} (x-a)^{n-j} (y-c)^j D^n f(a,c) \\ + \frac{1}{n!} \int_{c}^{y} \int_{a}^{x} (x-t)^n (y-s)^n \, d_t d_s \left(D^n f(t,s)\right), \end{aligned} \tag{2.5}$$

$$\begin{aligned} f(x,y) = \sum_{j=0}^{n} \frac{(-1)^j}{j!} \binom{n}{j} (x-a)^{n-j} (d-y)^j D^n f(a,d) \\ + \frac{(-1)^{n+1}}{n!} \int_{y}^{d} \int_{a}^{x} (x-t)^n (s-y)^n \, d_t d_s \left(D^n f(t,s)\right), \end{aligned} \tag{2.6}$$

$$\begin{aligned} f(x,y) = \sum_{j=0}^{n} \frac{(-1)^j}{j!} \binom{n}{j} (b-x)^{n-j} (y-c)^j D^n f(b,c) \\ + \frac{(-1)^{n+1}}{n!} \int_{c}^{y} \int_{x}^{b} (t-x)^n (y-s)^n \, d_t d_s \left(D^n f(t,s)\right) \end{aligned} \tag{2.7}$$

$$\begin{aligned} f(x,y) = \sum_{j=0}^{n} \frac{1}{j!} \binom{n}{j} (b-x)^{n-j} (d-y)^j D^n f(b,d) \\ + \frac{1}{n!} \int_{y}^{d} \int_{x}^{b} (t-x)^n (s-y)^n \, d_t d_s \left(D^n f(t,s)\right) \end{aligned} \tag{2.8}$$

for ny $(x,y) \in Q$.

Now, by multiplying (2.5) with $(b-x)(d-y)$, (2.6) with $(b-x)(y-c)$, (2.7) with $(x-a)(d-y)$, (2.8) with $(x-a)(y-c)$, we get

$$
\begin{aligned}
&(2.9)\quad (b-x)(d-y)f(x,y)\\
&=(b-x)(d-y)f(a,c)+(b-x)(d-y)\sum_{j=1}^{n}\frac{1}{j!}\left(\begin{array}{c} n \\ j \end{array}\right)(x-a)^{n-j}(y-c)^{j}D^{n}f(a,c)\\
&\qquad+\frac{1}{n!}(b-x)(d-y)\int_{c}^{y}\int_{a}^{x}(x-t)^{n}(y-s)^{n}d_{t}d_{s}(D^{n}f(t,s)),
\end{aligned}
$$

$$
\begin{aligned}
&(2.10)\quad (b-x)(y-c)f(x,y)\\
&=(b-x)(y-c)f(a,d)+(b-x)(y-c)\sum_{j=0}^{n}\frac{(-1)^{j}}{j!}\left(\begin{array}{c} n \\ j \end{array}\right)(x-a)^{n-j}(d-y)^{j}D^{n}f(a,d)\\
&\qquad+\frac{(-1)^{n+1}}{n!}(b-x)(y-c)\int_{y}^{d}\int_{a}^{x}(x-t)^{n}(s-y)^{n}d_{t}d_{s}(D^{n}f(t,s)),
\end{aligned}
$$

$$
\begin{aligned}
&(2.11)\quad (x-a)(d-y)f(x,y)\\
&=(x-a)(d-y)f(b,c)+(x-a)(d-y)\sum_{j=0}^{n}\frac{(-1)^{j}}{j!}\left(\begin{array}{c} n \\ j \end{array}\right)(b-x)^{n-j}(y-c)^{j}D^{n}f(b,c)\\
&\qquad+\frac{(-1)^{n+1}}{n!}(x-a)(d-y)\int_{c}^{y}\int_{x}^{b}(t-x)^{n}(y-s)^{n}d_{t}d_{s}(D^{n}f(t,s))
\end{aligned}
$$

$$
\begin{aligned}
&(2.12)\quad (x-a)(y-c)f(x,y)\\
&=(x-a)(y-c)f(b,d)+(x-a)(y-c)\sum_{j=0}^{n}\frac{1}{j!}\left(\begin{array}{c} n \\ j \end{array}\right)(b-x)^{n-j}(d-y)^{j}D^{n}f(b,d)\\
&\qquad+\frac{1}{n!}(x-a)(y-c)\int_{y}^{d}\int_{x}^{b}(t-x)^{n}(s-y)^{n}d_{t}d_{s}(D^{n}f(t,s))
\end{aligned}
$$

respectively, for ny $(x,y)\in Q$.

Finally, by adding the equalities (2.9)–(2.12) and dividing the sum with $(b-a)(d-c)$, we obtain

$$\begin{aligned}
f(x,y) = {} & \frac{1}{(b-a)(d-c)}\left[(b-x)(d-y)f(a,c)+(b-x)(y-c)f(a,d)\right.\\
& \left.+(x-a)(d-y)f(b,c)+(x-a)(y-c)f(b,d)\right]\\
& +\frac{(y-c)(d-y)}{(b-a)(d-c)}\sum_{j=1}^{n}\frac{1}{j!}\binom{n}{j}\left\{(b-x)(x-a)^{n-j}\left[(y-c)^{j-1}D^{n}f(a,c)\right.\right.\\
& \left.+(-1)^{j}(d-y)^{j-1}D^{n}f(a,d)\right]+(x-a)(b-x)^{n-j}\left[(-1)^{j}(y-c)^{j-1}D^{n}f(b,c)\right.\\
& \left.\left.+(d-y)^{j-1}D^{n}f(b,d)\right]\right\}\\
& +\frac{1}{n!}\frac{(b-x)(d-y)}{(b-a)(d-c)}\int_{c}^{y}\int_{a}^{x}(x-t)^{n}(y-s)^{n}\,d_{t}d_{s}\left(D^{n}f(t,s)\right)\\
& +\frac{(-1)^{n+1}}{n!}\frac{(b-x)(y-c)}{(b-a)(d-c)}\int_{y}^{d}\int_{a}^{x}(x-t)^{n}(s-y)^{n}\,d_{t}d_{s}\left(D^{n}f(t,s)\right),\\
& +\frac{(-1)^{n+1}}{n!}\frac{(x-a)(d-y)}{(b-a)(d-c)}\int_{c}^{y}\int_{x}^{b}(t-x)^{n}(y-s)^{n}\,d_{t}d_{s}\left(D^{n}f(t,s)\right)\\
& +\frac{1}{n!}\frac{(x-a)(y-c)}{(b-a)(d-c)}\int_{y}^{d}\int_{x}^{b}(t-x)^{n}(s-y)^{n}\,d_{t}d_{s}\left(D^{n}f(t,s)\right)
\end{aligned}$$

which gives the desired representation (2.1). □

**Remark 1.** *The case $n=0$ provides the representation*

$$\begin{aligned}
f(x,y) = {} & \frac{1}{(b-a)(d-c)}\left[(b-x)(d-y)f(a,c)+(b-x)(y-c)f(a,d)\right.\\
& \left.+(x-a)(d-y)f(b,c)+(x-a)(y-c)f(b,d)\right]\\
& +\frac{1}{(b-a)(d-c)}\int_{c}^{d}\int_{a}^{b}S_{0}(x,t;y,s)\,d_{t}d_{s}\left(f(t,s)\right)
\end{aligned}\tag{2.13}$$

*for any $x\in Q$, where*

$$S_{0}(x,t;y,s)=\begin{cases}(b-x)(d-y), & a\le t\le x, \quad c\le s\le y\\ (x-a)(d-y), & x<t\le b, \quad c\le s\le y\\ (b-x)(y-c), & a\le t\le x, \quad y<s\le d\\ (x-a)(y-c), & x<t\le b, \quad y<s\le d\end{cases}$$

*and $f$ is of bounded variation on $Q$.*

The case when $n=0$, provides an integral formula which compare any value of $f(x,y)$, $(x,y)\in Q$ with the values of the function and its derivatives at the rectangle end points (the corners of the rectangle generated by the end points). More specific, the Rectangular Mid-point value of $f$ can be represented in the following corollary:

**Corollary 1.** *With the assumptions of Theorem 3 for $f$ and $Q$, we have the identity*

$$
\begin{aligned}
(2.14)\quad f\left(\frac{a+b}{2},\frac{c+d}{2}\right) &= \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4} \\
&+\frac{1}{2^{n+2}}\sum_{j=1}^{n}\frac{1}{j!}\left(\begin{array}{c} n \\ j \end{array}\right)(b-a)^{n-j}(d-c)^{j}\left\{D^{n}f(a,c)+(-1)^{j}D^{n}f(a,d)\right. \\
&\left.+(-1)^{j}D^{n}f(b,c)+D^{n}f(b,d)\right\} \\
&+\frac{1}{(b-a)(d-c)}\int_{c}^{d}\int_{a}^{b}M_{n}(t,s)\,d_{t}d_{s}\left(D^{n}f(t,s)\right)
\end{aligned}
$$

*where,* $D^{n}f(t,s)=\frac{\partial^{n}f}{\partial t^{n-j}\partial s^{j}}(t,s)$ *and*

$$
M_{n}(t,s)=\frac{(b-a)(d-c)}{4n!}\left\{\begin{array}{ll}
\left(\frac{a+b}{2}-t\right)^{n}\left(\frac{c+d}{2}-s\right)^{n}, & a\le t\le\frac{a+b}{2},\quad c\le s\le\frac{c+d}{2} \\
(-1)^{n}\left(t-\frac{a+b}{2}\right)^{n}\left(\frac{c+d}{2}-s\right)^{n}, & \frac{a+b}{2}<t\le b,\quad c\le s\le\frac{c+d}{2} \\
(-1)^{n}\left(\frac{a+b}{2}-t\right)^{n}\left(s-\frac{c+d}{2}\right)^{n}, & a\le t\le\frac{a+b}{2},\quad \frac{c+d}{2}<s\le d \\
\left(t-\frac{a+b}{2}\right)^{n}\left(s-\frac{c+d}{2}\right)^{n}, & \frac{a+b}{2}<t\le b,\quad \frac{c+d}{2}<s\le d
\end{array}\right.
$$

On utilizing the following notations

$$
\begin{aligned}
(2.15)\quad \mathcal{A}(f,Q) &= \frac{1}{(b-a)(d-c)}\left[(b-x)(d-y)f(a,c)+(b-x)(y-c)f(a,d)\right. \\
&\left.+(x-a)(d-y)f(b,c)+(x-a)(y-c)f(b,d)\right]+\frac{(y-c)(d-y)}{(b-a)(d-c)} \\
&\times\sum_{j=1}^{n}\frac{1}{j!}\left(\begin{array}{c} n \\ j \end{array}\right)\left\{(b-x)(x-a)^{n-j}\left[(y-c)^{j-1}D^{n}f(a,c)+(-1)^{j}(d-y)^{j-1}D^{n}f(a,d)\right]\right. \\
&\left.+(x-a)(b-x)^{n-j}\left[(-1)^{j}(y-c)^{j-1}D^{n}f(b,c)+(d-y)^{j-1}D^{n}f(b,d)\right]\right\}
\end{aligned}
$$

and

$$
(2.16)\qquad \mathcal{B}_{n}(f,Q):=\frac{1}{(b-a)(d-c)}\int_{c}^{d}\int_{a}^{b}S_{n}(x,t;y,s)\,d_{t}d_{s}\left(D^{n}f(t,s)\right)
$$

under the assumptions of Theorem 3, we can approximate the function $f$ utilizing the polynomials $\mathcal{A}_{n}(f,Q)$ with the error $\mathcal{B}_{n}(f,Q)$ . In other words, we have

$$
f(x,y)=\mathcal{A}_{n}(f,Q)+\mathcal{B}_{n}(f,Q)
$$

for any $(x,y)\in Q$ .

The error $\mathcal{B}_{n}(f,Q)$ of a bounded bivariation function $f(x,y)$ satisfies the following bounds:

**Theorem 4.** *With the assumptions of Theorem 3 for $f$ and $Q$, we have*

$$
\begin{aligned}
(2.17)\qquad & |\mathcal{B}_n(f,Q)| \\
& \leq \frac{1}{n!\,(b-a)\,(d-c)} \cdot \max\left\{(y-c)^n\,(d-y),\,(y-c)\,(d-y)^n\right\} \\
& \qquad \times \max\left\{(x-a)^n\,(b-x),\,(x-a)\,(b-x)^n\right\} \cdot \bigvee_c^d \bigvee_a^b (D^n f) \\
& \leq \frac{(b-a)^n\,(d-c)^n}{2^{2n+2}(n)!} \cdot \bigvee_c^d \bigvee_a^b (D^n f)
\end{aligned}
$$

*Proof.* Using the inequality for the Riemann–Stieltjes integral of continuous integrands and bounded bivariation integrators, we have

$$
\begin{aligned}
|\mathcal{B}_n(f,Q)| = \Bigg| & \frac{1}{n!\,(b-a)\,(d-c)} \left[ \int_c^y \int_a^x (x-t)^n\,(b-x)\,(y-s)^n\,(d-y) d_t d_s\,(D^n f(t,s)) \right. \\
& + \int_y^d \int_a^x (-1)^{n+1}\,(x-t)^n\,(b-x)\,(s-y)^n\,(y-c) d_t d_s\,(D^n f(t,s)) \\
& + \int_c^y \int_x^b (-1)^{n+1}\,(t-x)^n\,(x-a)\,(y-s)^n\,(d-y) d_t d_s\,(D^n f(t,s)) \\
& \left. + \int_y^d \int_x^b (t-x)^n\,(x-a)\,(y-s)^n\,(d-y) d_t d_s\,(D^n f(t,s)) \right] \Bigg| \\
\leq & \frac{1}{n!\,(b-a)\,(d-c)} \left[ \left| \int_c^y \int_a^x (x-t)^n\,(b-x)\,(y-s)^n\,(d-y) d_t d_s\,(D^n f(t,s)) \right| \right. \\
& + \left| \int_y^d \int_a^x (-1)^{n+1}\,(x-t)^n\,(b-x)\,(s-y)^n\,(y-c) d_t d_s\,(D^n f(t,s)) \right| \\
& + \left| \int_c^y \int_x^b (-1)^{n+1}\,(t-x)^n\,(x-a)\,(y-s)^n\,(d-y) d_t d_s\,(D^n f(t,s)) \right| \\
& \left. + \left| \int_y^d \int_x^b (t-x)^n\,(x-a)\,(y-s)^n\,(d-y) d_t d_s\,(D^n f(t,s)) \right| \right]
\end{aligned}
$$

$$
\begin{aligned}
&\leq \frac{1}{n!\,(b-a)\,(d-c)}\Bigg[(b-x)\,(d-y)\max_{\substack{t\in[a,x]\\ s\in[c,y]}}\left\{(x-t)^n\,(y-s)^n\right\}\cdot\bigvee_c^y\bigvee_a^x\left(D^nf\right)\\
&\quad+(b-x)\,(y-c)\max_{\substack{t\in[a,x]\\ s\in[y,d]}}\left\{(x-t)^n\,(s-y)^n\right\}\cdot\bigvee_y^d\bigvee_a^x\left(D^nf\right)\\
&\quad+(x-a)\,(d-y)\max_{\substack{t\in[x,b]\\ s\in[c,y]}}\left\{(t-x)^n\,(y-s)^n\right\}\cdot\bigvee_c^y\bigvee_x^b\left(D^nf\right)\\
&\quad+(x-a)\,(y-c)\max_{\substack{t\in[x,b]\\ s\in[y,d]}}\left\{(t-x)^n\,(s-y)^n\right\}\cdot\bigvee_y^d\bigvee_x^b\left(D^nf\right)\Bigg]\\
&\leq \frac{1}{n!\,(b-a)\,(d-c)}\Bigg[(x-a)^n\,(b-x)\,(y-c)^n\,(d-y)\cdot\bigvee_c^y\bigvee_a^x\left(D^nf\right)\\
&\quad+(x-a)^n\,(b-x)\,(d-y)^n\,(y-c)\cdot\bigvee_y^d\bigvee_a^x\left(D^nf\right)\\
&\quad+(b-x)^n\,(x-a)\,(y-c)^n\,(d-y)\cdot\bigvee_c^y\bigvee_x^b\left(D^nf\right)\\
&\quad+(b-x)^n\,(x-a)\,(d-y)^n\,(y-c)\cdot\bigvee_y^d\bigvee_x^b\left(D^nf\right)\Bigg]\\
&\leq \frac{1}{n!\,(b-a)\,(d-c)}\Bigg[(x-a)^n\,(b-x)\max\left\{(y-c)^n\,(d-y),(y-c)\,(d-y)^n\right\}\cdot\bigvee_c^d\bigvee_a^x\left(D^nf\right)\\
&\quad+(x-a)\,(b-x)^n\max\left\{(y-c)^n\,(d-y),(y-c)\,(d-y)^n\right\}\cdot\bigvee_c^d\bigvee_x^b\left(D^nf\right)\Bigg]\\
&\leq \frac{1}{n!\,(b-a)\,(d-c)}\cdot\max\left\{(y-c)^n\,(d-y),(y-c)\,(d-y)^n\right\}\\
&\quad\times\max\left\{(x-a)^n\,(b-x),(x-a)\,(b-x)^n\right\}\cdot\bigvee_c^d\bigvee_a^b\left(D^nf\right)\\
&\leq \frac{(b-a)^n\,(d-c)^n}{2^{2n+2}(n)!}\cdot\bigvee_c^d\bigvee_a^b\left(D^nf\right),
\end{aligned}
$$

and the first inequality in (2.17) is proved. □

**Remark 2.** *Under the assumptions of Theorem 4, for all $(x,y)\in Q$, the case $n=0$ provides the following inequality:*

$$\begin{aligned}\left|\mathcal{B}_0(f,Q)\right| &\le \left[\frac{1}{2}+\frac{\left|x-\frac{a+b}{2}\right|}{b-a}\right]\left[\frac{1}{2}+\frac{\left|y-\frac{c+d}{2}\right|}{d-c}\right]\cdot\bigvee_c^d\bigvee_a^b(f)\\ &\le \frac{1}{4}\cdot\bigvee_c^d\bigvee_a^b(f).\end{aligned}$$

• **Midpoint formula:** Now, if we denote

$$\begin{aligned}\mathcal{E}_M^n(f;Q) &= \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\\ &+\frac{1}{2^{n+2}}\sum_{j=1}^{n}\frac{1}{j!}\binom{n}{j}(b-a)^{n-j}(d-c)^j\left\{D^nf(a,c)+(-1)^jD^nf(a,d)\right.\\ &\qquad\left.+(-1)^jD^nf(b,c)+D^nf(b,d)\right\}\end{aligned}\tag{2.18}$$

and

$$\mathcal{F}_M^n(f;Q)=\frac{1}{(b-a)(d-c)}\int_c^d\int_a^b M_n(t,s)\,d_td_s\left(D^nf(t,s)\right)\tag{2.19}$$

where,

$$M_n(t,s)=\frac{(b-a)(d-c)}{4n!}\times\begin{cases}\left(\frac{a+b}{2}-t\right)^n\left(\frac{c+d}{2}-s\right)^n, & a\le t\le\frac{a+b}{2},\quad c\le s\le\frac{c+d}{2}\\ (-1)^{n+1}\left(t-\frac{a+b}{2}\right)^n\left(\frac{c+d}{2}-s\right)^n, & \frac{a+b}{2}<t\le b,\quad c\le s\le\frac{c+d}{2}\\ (-1)^{n+1}\left(\frac{a+b}{2}-t\right)^n\left(s-\frac{c+d}{2}\right)^n, & a\le t\le\frac{a+b}{2},\quad\frac{c+d}{2}<s\le d\\ \left(t-\frac{a+b}{2}\right)^n\left(s-\frac{c+d}{2}\right)^n, & \frac{a+b}{2}<t\le b,\quad\frac{c+d}{2}<s\le d\end{cases}$$

then we can approximate the value of a function at its midpoint in terms of values of the function and its partial derivatives taken at the end points with error $\mathcal{F}_M(f;Q)$. Namely, we have the representation formula

$$f\left(\frac{a+b}{2},\frac{c+d}{2}\right)=\mathcal{E}_M^n(f;Q)+\mathcal{F}_M^n(f;Q)\tag{2.20}$$

The absolute value of the error can be bounded as follows:

**Corollary 2.** *With the assumptions of Theorem 4 for $f$, $Q$ and $n$, we have the inequality*

$$\left|\mathcal{F}_M^n(f;Q)\right|\le\frac{(b-a)^n(d-c)^n}{2^{2n+2}n!}\cdot\bigvee_c^d\bigvee_a^b(f)\tag{2.21}$$

For another assumptions on $f$, the case when $D^nf$ are absolutely continuous on $Q$ for all $(n\ge 0)$, we give the following estimates for the remainder $\mathcal{B}_n(f,Q)$:

**Theorem 5.** *Let $f:Q\to\mathbb{R}$ be a real valued function which has continuous $(n-1)$ partial derivatives $(n\ge 1)$. If the $n$-th partial derivatives $D^nf$ $(n\ge 1)$ are*

*absolutely continuous on $Q$, then for any $(x,y)\in Q$ we have*

(2.22)
$$|\mathcal{B}_n(f,Q)|\leq\begin{cases}\frac{1}{(n)!(n+1)^2(b-a)(d-c)}\left[(b-x)(x-a)^{n+1}+(x-a)(b-x)^{n+1}\right]\\ \times\left[(d-y)(y-c)^{n+1}+(y-c)(d-y)^{n+1}\right]\cdot\left\|D^{n+1}f\right\|_{Q,\infty}, & \text{if } D^{n+1}f\in L_\infty(Q);\\ \frac{1}{(n)!(nq+1)^{1/q}(b-a)(d-c)}\left[(b-x)(x-a)^{n+1/q}+(x-a)(b-x)^{n+1/q}\right]\\ \times\left[(d-y)(y-c)^{n+1/q}+(y-c)(d-y)^{n+1/q}\right]\cdot\left\|D^{n+1}f\right\|_{Q,p}, & \text{if } D^{n+1}f\in L_p(Q),\\ & p>1,\frac{1}{p}+\frac{1}{q}=1;\end{cases}$$

*Proof.* Since $D^nf$ is absolutely continuous on $Q$ then for any $(x,y)\in Q$ we have the representation

$$\begin{aligned}f(x,y)&=\frac{1}{(b-a)(d-c)}\left[(b-x)(d-y)f(a,c)+(b-x)(y-c)f(a,d)\right.\\&\left.+(x-a)(d-y)f(b,c)+(x-a)(y-c)f(b,d)\right]+\frac{(y-c)(d-y)}{(b-a)(d-c)}\\&\times\sum_{j=1}^{n}\frac{1}{j!}\binom{n}{j}\left\{(b-x)(x-a)^{n-j}\left[(y-c)^{j-1}D^nf(a,c)+(-1)^j(d-y)^{j-1}D^nf(a,d)\right]\right.\\&\left.+(x-a)(b-x)^{n-j}\left[(-1)^j(y-c)^{j-1}D^nf(b,c)+(d-y)^{j-1}D^nf(b,d)\right]\right\}\\&+\frac{1}{(b-a)(d-c)}\int_c^d\int_a^b S_n(x,t;y,s)D^{n+1}f(t,s)\,dtds,\end{aligned}\tag{2.23}$$

where the integral is considered in the Lebesgue sense and the kernel $S_n(x,t;y,s)$ is given in Theorem 3. Utilizing the properties of the Stieltjes integral, we have

$$\begin{aligned}|\mathcal{B}_n(f;Q)|&=\frac{1}{(b-a)(d-c)}\left|\int_c^d\int_a^b S_n(x,t;y,s)D^{n+1}f(t,s)\,dtds\right|\\&=\left|\frac{1}{n!(b-a)(d-c)}\left[\int_c^y\int_a^x(x-t)^n(b-x)(y-s)^n(d-y)D^{n+1}f(t,s)\,dtds\right.\right.\\&+\int_y^d\int_a^x(-1)^{n+1}(x-t)^n(b-x)(s-y)^n(y-c)D^{n+1}f(t,s)\,dtds\\&+\int_c^y\int_x^b(-1)^{n+1}(t-x)^n(x-a)(y-s)^n(d-y)D^{n+1}f(t,s)\,dtds\\&\left.\left.+\int_y^d\int_x^b(t-x)^n(x-a)(y-s)^n(d-y)D^{n+1}f(t,s)\,dtds\right]\right|\end{aligned}$$

$$
\begin{aligned}
&\leq \frac{1}{n!\,(b-a)\,(d-c)}\left[\left|\int_c^y\int_a^x (x-t)^n\,(b-x)\,(y-s)^n\,(d-y)\,D^{n+1}f\,(t,s)\,dtds\right|\right.\\
&\quad+\left|\int_y^d\int_a^x (-1)^{n+1}\,(x-t)^n\,(b-x)\,(s-y)^n\,(y-c)\,D^{n+1}f\,(t,s)\,dtds\right|\\
&\quad+\left|\int_c^y\int_x^b (-1)^{n+1}\,(t-x)^n\,(x-a)\,(y-s)^n\,(d-y)\,D^{n+1}f\,(t,s)\,dtds\right|\\
&\quad+\left.\left|\int_y^d\int_x^b (t-x)^n\,(x-a)\,(y-s)^n\,(d-y)\,D^{n+1}f\,(t,s)\,dtds\right|\right]\\
&\leq \frac{1}{n!\,(b-a)\,(d-c)}\left[(b-x)\,(d-y)\int_c^y\int_a^x (x-t)^n\,(y-s)^n\left|D^{n+1}f\,(t,s)\right|dtds\right.\\
&\quad+(y-c)\,(b-x)\int_y^d\int_a^x (x-t)^n\,(s-y)^n\left|D^{n+1}f\,(t,s)\right|dtds\\
&\quad+(d-y)\,(x-a)\int_c^y\int_x^b (t-x)^n\,(y-s)^n\left|D^{n+1}f\,(t,s)\right|dtds\\
&\quad+\left.(d-y)\,(x-a)\int_y^d\int_x^b (t-x)^n\,(y-s)^n\left|D^{n+1}f\,(t,s)\right|dtds\right].
\end{aligned}
$$

Utilizing properties of integration together with Hölder integral inequality for the Lebesgue integral we have

$$
\begin{aligned}
&\int_c^y\int_a^x (x-t)^n\,(y-s)^n\left|D^{n+1}f\,(t,s)\right|dtds \qquad (2.24)\\
&=\begin{cases}
\frac{(x-a)^{n+1}(y-c)^{n+1}}{(n+1)^2}\cdot\left\|D^{n+1}f\right\|_{[a,x]\times[c,y],\infty}, & D^{n+1}f\in L_\infty[a,x]\times[c,y];\\
\frac{(x-a)^{n+1/q}(y-c)^{n+1/q}}{(nq+1)^{1/q}}\cdot\left\|D^{n+1}f\right\|_{[a,x]\times[c,y],p}, & D^{n+1}f\in L_p[a,x]\times[c,y],\\
& p>1,\frac{1}{p}+\frac{1}{q}=1;
\end{cases}
\end{aligned}
$$

$$
\begin{aligned}
&\int_y^d\int_a^x (x-t)^n\,(s-y)^n\left|D^{n+1}f\,(t,s)\right|dtds \qquad (2.25)\\
&=\begin{cases}
\frac{(x-a)^{n+1}(d-y)^{n+1}}{(n+1)^2}\cdot\left\|D^{n+1}f\right\|_{[a,x]\times[y,d],\infty}, & D^{n+1}f\in L_\infty[a,x]\times[y,d];\\
\frac{(x-a)^{n+1/q}(d-y)^{n+1/q}}{(nq+1)^{1/q}}\cdot\left\|D^{n+1}f\right\|_{[a,x]\times[y,d],p}, & D^{n+1}f\in L_p[a,x]\times[y,d],\\
& p>1,\frac{1}{p}+\frac{1}{q}=1;
\end{cases}
\end{aligned}
$$

$$
\begin{aligned}
&\int_c^y\int_x^b (t-x)^n\,(y-s)^n\left|D^{n+1}f\,(t,s)\right|dtds \qquad (2.26)\\
&=\begin{cases}
\frac{(b-x)^{n+1}(y-c)^{n+1}}{(n+1)^2}\cdot\left\|D^{n+1}f\right\|_{[x,b]\times[c,y],\infty}, & D^{n+1}f\in L_\infty[x,b]\times[c,y];\\
\frac{(b-x)^{n+1/q}(y-c)^{n+1/q}}{(nq+1)^{1/q}}\cdot\left\|D^{n+1}f\right\|_{[x,b]\times[c,y],p}, & D^{n+1}f\in L_p[x,b]\times[c,y],\\
& p>1,\frac{1}{p}+\frac{1}{q}=1;
\end{cases}
\end{aligned}
$$

and

$$\int_y^d \int_x^b (t-x)^n (y-s)^n \left|D^{n+1} f(t,s)\right| dtds \tag{2.27}$$

$$= \begin{cases} \frac{(b-x)^{n+1}(d-y)^{n+1}}{(n+1)^2} \cdot \left\|D^{n+1} f\right\|_{[x,b]\times[y,d],\infty}, & D^{n+1} f \in L_\infty [x,b]\times[y,d]; \\ \frac{(b-x)^{n+1/q}(d-y)^{n+1/q}}{(nq+1)^{1/q}} \cdot \left\|D^{n+1} f\right\|_{[x,b]\times[y,d],p}, & D^{n+1} f \in L_p [x,b]\times[y,d], \\ & p>1, \frac{1}{p}+\frac{1}{q}=1; \end{cases}$$

Adding (2.24)–(2.27) we deduce that

$$\begin{aligned} &|\mathcal{B}_n (f,Q)| \\ &\leq \frac{(b-x)(d-y)}{n!(b-a)(d-c)} \times \begin{cases} \frac{(x-a)^{n+1}(y-c)^{n+1}}{(n+1)^2} \cdot \left\|D^{n+1} f\right\|_{[a,x]\times[c,y],\infty}, & D^{n+1} f \in L_\infty [a,x]\times[c,y]; \\ \frac{(x-a)^{n+1/q}(y-c)^{n+1/q}}{(nq+1)^{1/q}} \cdot \left\|D^{n+1} f\right\|_{[a,x]\times[c,y],p}, & D^{n+1} f \in L_p [a,x]\times[c,y], \\ & p>1, \frac{1}{p}+\frac{1}{q}=1; \end{cases} \\ &+\frac{(b-x)(y-c)}{n!(b-a)(d-c)} \times \begin{cases} \frac{(x-a)^{n+1}(d-y)^{n+1}}{(n+1)^2} \cdot \left\|D^{n+1} f\right\|_{[a,x]\times[y,d],\infty}, & D^{n+1} f \in L_\infty [a,x]\times[y,d]; \\ \frac{(x-a)^{n+1/q}(d-y)^{n+1/q}}{(nq+1)^{1/q}} \cdot \left\|D^{n+1} f\right\|_{[a,x]\times[y,d],p}, & D^{n+1} f \in L_p [a,x]\times[y,d], \\ & p>1, \frac{1}{p}+\frac{1}{q}=1; \end{cases} \\ &+\frac{(x-a)(d-y)}{n!(b-a)(d-c)} \times \begin{cases} \frac{(b-x)^{n+1}(y-c)^{n+1}}{(n+1)^2} \cdot \left\|D^{n+1} f\right\|_{[x,b]\times[c,y],\infty}, & D^{n+1} f \in L_\infty [x,b]\times[c,y]; \\ \frac{(b-x)^{n+1/q}(y-c)^{n+1/q}}{(nq+1)^{1/q}} \cdot \left\|D^{n+1} f\right\|_{[x,b]\times[c,y],p}, & D^{n+1} f \in L_p [x,b]\times[c,y], \\ & p>1, \frac{1}{p}+\frac{1}{q}=1; \end{cases} \\ &+\frac{(x-a)(y-c)}{n!(b-a)(d-c)} \times \begin{cases} \frac{(b-x)^{n+1}(d-y)^{n+1}}{(n+1)^2} \cdot \left\|D^{n+1} f\right\|_{[x,b]\times[y,d],\infty}, & D^{n+1} f \in L_\infty [x,b]\times[y,d]; \\ \frac{(b-x)^{n+1/q}(d-y)^{n+1/q}}{(nq+1)^{1/q}} \cdot \left\|D^{n+1} f\right\|_{[x,b]\times[y,d],p}, & D^{n+1} f \in L_p [x,b]\times[y,d], \\ & p>1, \frac{1}{p}+\frac{1}{q}=1; \end{cases} \\ &\leq \begin{cases} \frac{1}{(n)!(n+1)^2(b-a)(d-c)} \left[(b-x)(x-a)^{n+1}+(x-a)(b-x)^{n+1}\right] \\ \times \left[(d-y)(y-c)^{n+1}+(y-c)(d-y)^{n+1}\right] \cdot \left\|D^{n+1} f\right\|_{Q,\infty}, \\ \qquad \text{if } D^{n+1} f \in L_\infty (Q_{x,c}^{b,y}); \\ \frac{1}{(n)!(nq+1)^{1/q}(b-a)(d-c)} \left[(b-x)(x-a)^{n+1/q}+(x-a)(b-x)^{n+1/q}\right] \\ \times \left[(d-y)(y-c)^{n+1/q}+(y-c)(d-y)^{n+1/q}\right] \cdot \left\|D^{n+1} f\right\|_{Q,p}, \\ \qquad \text{if } D^{n+1} f \in L_p (Q_{x,c}^{b,y}), \\ \qquad p>1, \frac{1}{p}+\frac{1}{q}=1; \end{cases} \end{aligned}$$

□

Thus, the value of the function $f$ can be approximated at its midpoint in terms of values of $f$ and its partial derivatives taken at the end points with error $\mathcal{F}_M(f;Q)$, i.e.,

$$f\left(\frac{a+b}{2},\frac{c+d}{2}\right)=\mathcal{E}_M\left(f;Q\right)+\mathcal{F}_M\left(f;Q\right),$$

such that the error $\mathcal{F}_M(f;Q)$ satisfies the following bounds:

**Corollary 3.** *With the assumptions of Theorem 5 for $f$, $Q$ and $n$, we have the inequality*

$$\left|\mathcal{F}_M^n\left(f;Q\right)\right|\leq\frac{1}{n!}\times\begin{cases}\frac{(b-a)^{n+1}(d-c)^{n+1}}{2^{2n+2}(n+1)^2}\cdot\left\|D^{n+1}f\right\|_{Q,\infty}, & \text{if } D^{n+1}f\in L_\infty(Q);\\ \frac{(b-a)^{n+1/q}(d-c)^{n+1/q}}{2^{2n+2/q}(nq+1)^{1/q}}\cdot\left\|D^{n+1}f\right\|_{Q,p}, & \text{if } D^{n+1}f\in L_p(Q),\\ & p>1,\frac{1}{p}+\frac{1}{q}=1;\end{cases}.$$

*In particular, for $n=0$, we have*

$$\left|\mathcal{F}_M^0\left(f;Q\right)\right|\leq\begin{cases}\frac{(b-a)(d-c)}{4}\cdot\left\|Df\right\|_{Q,\infty}, & \text{if } Df\in L_\infty(Q);\\ \frac{(b-a)^{1/q}(d-c)^{1/q}}{4^{1/q}}\cdot\left\|Df\right\|_{Q,p}, & \text{if } Df\in L_p(Q),\\ & p>1,\frac{1}{p}+\frac{1}{q}=1;\end{cases}.$$

**Remark 3.** *Different kinds of Hölder continuous functions of two variable have been defined in literature, for example a function $f:Q\to\mathbb{R}$ is to be of $(\beta_1,\beta_2)$–Hölder type mapping on the co-ordinate, if for all $(t_1,s_1),(t_1,s_1)\in Q$, there exist $H_1,H_2>0$ and $\beta_1,\beta_2>0$ such that*

$$\left|f\left(t_1,s_1\right)-f\left(t_2,s_2\right)\right|\leq H_1\left|t_1-t_2\right|^{\beta_1}+H_2\left|s_1-s_2\right|^{\beta_2}.$$

*If $\beta_1=\beta_2=1$, then $f$ is called $(L_1,L_2)$–Lipschitz on the co-ordinate, i.e.,*

$$\left|f\left(t_1,s_1\right)-f\left(t_2,s_2\right)\right|\leq L_1\left|t_1-t_2\right|+L_2\left|s_1-s_2\right|.$$

*Another definition of Hölder continuous mapping can be stated as:*

$$\left|f\left(t_1,s_1\right)-f\left(t_2,s_2\right)\right|\leq M_{\beta_1,\beta_2}\left(t_1,t_2;s_1,s_2\right).$$

*where, $M_{\beta_1,\beta_2}\left(t_1,t_2;s_1,s_2\right)$ is any real norm. In particular, if $\beta_1=\beta_2=1$, the usual Euclidean norm is well known as:*

$$M_{1,1}\left(t_1,t_2;s_1,s_2\right):=\sqrt{\left(t_1-s_1\right)^2+\left(t_2-s_2\right)^2}.$$

*Under any type of Hölder continuity, one may has several bounds for the error bounds $\mathcal{B}_n\left(f,Q\right)$ and $\mathcal{F}_M^n\left(f,Q\right)$ of the function $f$. We leave this to the interested reader (see* [1]*).*

## References


[1] M. W. Alomari, Numerous approximations of Riemann-Stieltjes double integrals, monograph 150 pages, 2009. Avaliable at `https://www.scribd.com/document/324214887`

[2] M. W. Alomari, New Grüss type inequalities for double integrals, *Appl. Math. Comp.*, 228 (2014) 102–107.

[3] G.A. Anastassiou, Univariate Ostrowski inequalities, *Monash. Math.,* **135** (3) (2002) 175–189. Revisited.

[4] N.S. Barnett and S.S. Dragomir, An Ostrowski type inequality for double integrals and applications for cubature formulae, *Soochow J. Math.* **27** (2001), 1–10.

[5] J.A. Clarkson, On double Riemann–Stieltjes integrals, *Bull. Amer. Math. Soc.* **39** (1933), 929–936.
[6] J.A. Clarkson and C. R. Adams, On definitions of bounded variation for functions of two variables, *Bull. Amer. Math. Soc.* **35** (1933), 824–854.
[7] P. Cerone, S.S. Dragomir and C.E.M. Pearce, A generalised trapezoid inequality for functions of bounded variation, *Turk. J. Math.*, 24 (2000), 147–163.
[8] S.S. Dragomir, Sharp bounds for the deviation of a function from the chord generated by its extremities and applications, *RGMIA Res. Rep. Coll.,* **10** (Suppl.) (2007) Preprint, Article 17 [Online http://rgmia.vu.edu.au/v10(E).html].
[9] S.S. Dragomir, Approximating real functions which possess $n$-th derivatives of bounded variation and applications, *Computers and Mathematics with Applications,* **56** (2008), 2268–2278.
[10] S.S. Dragomir, P. Cerone, N.S. Barnett and J. Roumeliotis, An inequality of the Ostrowski type for double integrals and applications for cubature formulae, *Tamsui Oxf. J. Math. Sci.* **16** (2000), 1–16.
[11] S.S. Dragomir, N.S. Barnett and P. Cerone, An Ostrowski type inequality for double integrals in terms of $L_p$-norms and applications in numerical integration, *Rev. Anal. Numer. Theor. Approx.* **32** (2003), 161–169.
[12] S.S. Dragomir and T.M. Rassias, Ostrowski type inequalities and applications in numerical integration, Boston: Kluwer Academic (2002).
[13] M. Fréchet, Extension au cas des intégrals multiples d'une définition de l'intégrale due à Stieltjes, *Nouvelles Annales de Mathématiques* **10** (1910), 241–256.
[14] S. Ghorpade and B. Limaye, A course in multivariable calculus and analysis, Springer: New York (2009).
[15] G. Hanna, Cubature rules from a generalized Taylor prespective, PH.D. Thesis, Victoria University (2009).
[16] J.E. Pečarić and A. Vukelić, Montgomery's identities for function of two variables, *J. Math. Anal. Appl.* **332** (2007), 617–630.
[17] Z. Liu, A generalization of two sharp inequalities in two independent variables, *Comp. Math. Appl.* **59** (2010) 2809–2814.
[18] Lü Zhongxue, On sharp inequalities of Simpson type and Ostrowski type in two independent variables, *Comp. Math. Appl.* **56** (2008) 2043–2047.

Department of Mathematics, Faculty of Science and Information Technology, Irbid National University, 2600 Irbid 21110, Jordan.
*E-mail address*: mwomath@gmail.com